Chapter

Standard Page

1

App. 3

Fish Products Standards and Methods Manual

Status

Date

Amend.no.11 12/10/2011

APPENDIX 3 CANADIAN GUIDELINES FOR CHEMICAL CONTAMINANTS AND TOXINS IN FISH AND FISH PRODUCTS

CONTAMINANTS	PRODUCT TYPE	ACTION LEVEL 1	
Mercury	All fish products (except swordfish, shark, fresh and frozen tuna, escolar, orange roughy and marlin)		
Mercury	Swordfish, shark, fresh and frozen tuna, escolar, orange roughy and marlin	1.0 ppm	
Arsenic	Fish protein concentrate	3.5 ppm	
Lead	Fish protein concentrate	0.5 ppm	
Fluoride	Fish protein concentrate	150 ppm	
2,3,7,8 TCDD (Dioxin)	All fish products	20 ppt *under review*	
DDT and Metabolites (DDD and DDE)	All fish products	5.0 ppm	
PCB	All fish products	2.0 ppm *under review*	
Piperonyl butoxide	Dried Cod	1.0 ppm	
Other agricultural chemicals or their derivatives	All fish products	0.1 ppm	

¹ Based on contaminants level of edible weight

Notes:

Sampling: Samples to consist of a minimum of 5 units representative of the lot. Analysis may be carried out on a composite of all sample units.

Criteria for action: A lot of fish will be considered reject if the sample value exceeds the action level. Fish or fish products exceeding these guidelines may be permitted for export if they do not violate regulations of the importing country.

TOXINS	PRODUCT TYPE	ACTION LEVEL
Histamine ' (Scombroid Poisoning)	Enzyme ripened products (e.g., anchovies, anchovy paste, fish sauce)	20 mg/100 g
Histamine ' (Scombroid Poisoning)	All other scombroid fish products (e.g., canned or fresh or frozen tuna, mackerel, mahi-mahi)	10 mg/100 g
Saxitoxins (PSP) 3	Molluscan shellfish (edible portion)	80 μg/100 g

<u>Chapter</u>

Standard

App. 3

<u>Page</u> -

2

Fish Products Standards

and Methods Manual

Status

<u>Date</u> Amend.no.11 12/10/2011

Domoic Acid (ASP)	Molluscan shellfish	(edible portion)	20 μg/g
Okadaic Acid (OA) + DTX1 + DTX2 + OA esters + DTX1 esters + DTX2 esters (DSP) ³	Molluscan shellfish	(edible portion)	0.2 μg/g* (interim)
Pectenotoxi ns: PTX-1, PTX-2, PTX- 3, PTX-4, PTX-6 and PTX-11	Molluscan shellfish	(edible portion)	0.2 μg/g

Chapter

Standard Page

App. 3

<u>Date</u>

3

Fish Products Standards and Methods Manual

Status

New

12/10/2011

ADDITIONAL COMMENTS:

² Histamine

- Samples are collected according to Sampling Plan 1 (AQL 6.5) for Initial inspection and Sampling Plan 2 (AQL 6.5) for reinspection.
- Any sample exceeding 50 mg/100 g will result in the lot being rejected with no right to reinspection.
- The acceptance number is that corresponding to the number for decomposition.
- ³ PSP, ASP and DSP (Paralytic Shellfish Poisoning, Amnesic Shellfish Poisoning Domoic Acid, Diarrhetic Shellfish Poisoning Okadaic Acid and/or DTX-1)
- Procedures for closure of shellfish areas, and possible recall of product due to samples of shellfish containing toxin levels equal to or greater than the above action levels can be found in Chapter 11 of the Canadian Shellfish Sanitation program
- The minimum acceptable sample is that which when shucked will produce 100 g of drained meats from 5 pooled sub-samples. Depending on the size of animals, the total number of shellfish required varies from 3 (geoduck) to 25 (pink scallops)

BACKGROUND LEVELS FOR NON-PERMITTED ADDITIVES

ADDITIVE ¹	PRODUCT TYPE	BACKGROUND LEVEL
Nitrites	All fish and fish products (except marine mammal meat ⁶)	15 ppm (see note 2)
Nitrates	All fish and fish products	15 ppm (see note 2)
Sulphites'	Clams (raw and canned)	10 ppm
Phosphates	Shrimp (raw, cooked and canned)	1.60 %
Phosphates	Scallops (raw)	1.47 %
Phosphates*	Fish fillets	1.37 %

Chapter

<u>Standar</u>d

12/10/2011

App. 3

New

Page

Fish Products Standards

Status Date and Methods Manual

Phosphates*	Crab (raw and cooked)	1.70 %
Phosphates	Lobster (raw and cooked)	1.47 %
Phosphates*	Surf clams (raw and cooked)	1.00 %

⁴ The compounds listed in this table are food additives; however some background levels may occur naturally in some foods.

Note:

- 1. If a processor can provide reliable data for naturally occurring background levels that are higher than those shown above, this may be considered before product action is taken.
- Some herbs, including parsley, contain high levels of naturally occurring 2. nitrates. This has to be considered when nitrates are detected in fish products containing herbs as an ingredient.

⁵ When the additive **is not** permitted, then the action level is the background level or detection limit; when the additive is permitted, then the action level is the background level or detection limit plus the permitted amount.

⁶ Marine mammals, including seals are included in the definition of "fish" as per the Canadian Food and Drug Regulations. Sodium nitrite is permitted in marine mammal meats at the maximum level of 200 ppm.

⁷ Calculated as sulphur dioxide.

⁸ Calculated as sodium phosphate, dibasic.